



## The Psychology of Polio as Prelude to Post-Polio Sequelae: Behavior Modification and Psychotherapy.

Drs. Nancy M. Frick, M.Div, LhD and Richard L. Bruno, HD, PhD

Orthopedics, 1991, 14(11): 1185-1193.

### ABSTRACT

Even as the physical causes and treatments for Post-Polio Sequelae (PPS) are being identified, psychological symptoms - chronic stress, anxiety, depression, compulsive and Type A behavior - are becoming evident in polio survivors. Importantly, these symptoms are not only themselves causing marked distress but also are preventing patients from making the lifestyle changes necessary to treat PPS. Neither clinicians nor polio survivors have paid sufficient attention to the acute polio experience, its conditioning of life-long patterns of behavior, its relationship to the development of PPS and its effect on the ability of polio survivors to cope with and treat their new symptoms. This paper describes the acute polio and post-polio experiences on the basis of patient histories, relates the experience of polio to the development of compulsive Type A behavior, links these behaviors to the physical and psychological symptoms reported in the National Post-Polio Surveys and presents a multimodal behavioral approach to the treatment of PPS by describing patients who have been treated by this Post-Polio Service.

*"I'll walk without crutches. I'll walk into a room without scaring everybody half to death. I'll stand easily enough in front of people so that they'll forget I'm a cripple." Franklin D. Roosevelt (1)*

**In the 1980's**, nearly half of the then 1.63 million American polio survivors reported new and unexpected symptoms - fatigue, muscle weakness and pain - that have come to be called Post-Polio Sequelae (PPS) (2,3). While the medical community focused its attention on these new physical symptoms, polio survivors' psychological reactions to PPS emerged. Fear, disbelief, denial and confusion were experienced first since PPS were completely unexpected (4). The 1985 National Survey of 738 polio survivors (5,6) revealed that 86% of the respondents never considered the possibility of developing new problems after having reached maximum recovery from polio, and 72% did not initially think that their new symptoms were related to having had polio. Since polio survivors did not know the cause of their new symptoms, they looked to the medical community for help.

Fear and confusion increased as medical ignorance and disinterest frustrated and angered polio survivors. Many physicians dismissed new symptoms as psychogenic, factitious or unrelated to polio. In the 1985 National Survey, 23% of respondents were told by physicians that the symptoms were "all in their minds" while 26% were told that the symptoms "could not possibly be related to having had polio." Other patients received frightening and unfounded diagnoses, such as "some kind of ALS" or recrudescence polio (3). Many patients were told that they would have to accept new symptoms and loss of function since all polio survivors were "just getting old" (4).

Now, in the second decade of PPS, the medical community has accepted the reality of PPS and is identifying effective treatments for new fatigue, weakness and pain. Even with these advances, additional psychological symptoms - chronic stress, anxiety, depression and compulsive, Type A behavior - are being identified in polio survivors. Importantly, these symptoms are not only themselves causing marked distress but also are preventing patients from making the lifestyle changes necessary to treat their PPS (6,7,8).

It appears that the factor central to the etiology of these psychological symptoms is that polio

survivors are being forced to cope with new symptoms and disability when many have not yet dealt with the emotional reality of their having had polio more than thirty years ago. Evidence for the postponement of coping with the acute polio experience was found in the 1985 National Survey, where 26% of respondents reported that they "did not think of themselves as having a disability" before developing PPS; with the onset of PPS 42% admitted to having a "second disability." (6)

Neither clinicians nor polio survivors have paid sufficient attention to the acute polio experience, its conditioning of life-long patterns of behavior, relationship to the development of PPS and its effect on individuals' ability to cope with and treat their new symptoms. It is the purpose of this paper to:

- 1) describe the acute polio and post-polio experiences on the basis of published reports and patient histories;
- 2) relate the experience of polio to the development of compulsive, Type A behavior;
- 3) link compulsive, Type A behavior to the physical and psychological symptoms reported in the National Post-Polio Surveys;
- 4) present a multimodal behavioral approach to the treatment of PPS by describing patients who have been treated by this Post-Polio Service during the past year.

**The Acute Polio Experience.** Polio survivors frequently report that the onset of PPS have forced them, often for the first time, to recall and examine their acute polio experience (9). With the onset of polio, these individuals understood that they had been stricken by the "Igreatly feared disease," believed that disability was likely and death was possible (10). These fears were amplified in those who knew others who had been disabled or even killed by polio (10). Polio survivors recall confusion and terror at being wrenched from their homes and being given by their parents into the care of strangers with whom they were required to live for weeks, months or even years (11). Even children too young to understand what was happening likely sensed their parents' fear and panic; they certainly felt the pain of their own illness and the desolation of being abandoned by their parents (11,12,13,14).

Individuals admitted to hospital quickly became aware of their complete loss of control and total dependency on the hospital staff. Many could not feed or toilet themselves; those with the most severe paralysis could not even move. Patients' were forced to rely totally on hospital staff if their most basic survival needs were to be met. If they were to survive, patients had little choice but to suppress their fears and follow their parents' parting instruction: "Be good, don't make trouble and do everything you're told" (11).

Separation from parents and loss of control were not the only sources of fear. Patients describe a hospital regimen of painful and often frightening therapies including boiling hot packs, splinting, bracing, stretching and exercise (15). Patients also report undergoing multiple surgeries, including muscle transfers, tendon lengthening and osteotomies. One especially brutal procedure that parents attempted to obtain for their children pulverized weakened muscles and their motor nerves with an electric riveter to promote axonal sprouting, a uniquely medieval attempt to prevent permanent visible disability (16,17). These procedures were administered usually without explanation and certainly without consent since, especially at the height of the epidemics, hospitals were overwhelmed by the sheer number of polio patients needing treatment (10,11,12,13,15,18,19). Excluding patients from treatment decisions and the failure to communicate information about their illness magnified the severity of the illness and often permitted patients inherent fear of death to emerge (14).

Questions or complaints about therapies were not infrequently met by staff anger or punishment (13,18). Staff anger was viewed as a very real threat to survival by patients who were isolated from

parents and totally dependent on the staff. One patient described this situation as placing her in "mortal danger" (15); she believed that the staff had to be placated if she were to survive. Patients discovered that staff anger could sometimes be prevented if they complied with the hospital regimen without questions. Full participation in therapies, without regard to pain or exhaustion, would sometimes garner brief attention or even praise. Thus, the expression of personal needs and emotion were punished by the staff while unquestioning compliance with and performance for the staff were reinforced.

Unfortunately, protection from punishment could not always be assured. Patients describe staff who acted with unnecessary cruelty in maintaining order and control on the ward. Family visits were restricted to only a few hours per week, reinforcing patients' feelings of parental abandonment and dependency upon staff (12,15, 20). Normal, child-like behaviors were punished excessively. Several patients report having been locked in completely dark closets overnight when they spoke or cried after "lights out." Even appropriate and necessary nursing care could be withheld for no apparent reason (15,18).

Many of the patients treated by The Post-Polio Institute have related other instances of psychological, physical and even sexual abuse at the hands of hospital staff. Instead of causing patients to complain, such abuse further reinforced patients' belief that survival depended upon the suppression of personal needs and emotion and unquestioning compliance with those in authority. This belief is evidenced in one patient's rationale for not reporting repeated sexual abuse: "I would have made my parents angry because I was not doing what I was told; the nurses, who I depended on for everything, would have punished me for making trouble for them. All I could do was stop feeling bad about it and smile."

Under these circumstances, some patients report having lost their sense of "identity" (18,20). Some report over identifying with the staff and even expressed the desire to become orthopedic surgeons when they grew up (21). Others apparently submitted so completely to the hospital staff that they did not wish to go home or even asked to return to the hospital following discharge (22,23).

It appears] that many polio survivors learned to deal with their abandonment, loss of control, fear, pain and abuse by submitting to those in authority, complying fully with external expectations and denying personal needs, physical and emotional pain and even their own individuality (24).

These behaviors are evident in this summary of the "Good Chart" written by polio wardmates at Baltimore's City Hospital (22):

LISTEN TO THE DOCTORS  
OBEY THE NURSES  
DO NOT FIGHT  
DO NOT BE BAD  
BE GOOD IN SCHOOL  
DO YOUR HOMEWORK  
DO NOT TALK AT DINNER or IN SCHOOL  
LITTLE FOLKS SHOULD BE SEEN AND NOT HEARD

When polio patients returned home as "polio survivors", the "Good Chart" became their prescription for proper behavior. They had learned very young and very well that submission, compliance and suppression of emotion were required if they were to escape "mortal danger"(15).

**Polio Survivors in the Community.** When polio survivors re-entered their communities, any special attention that may have been provided to them and their families during the acute polio often quickly stopped (22). Friends of polio survivors' families, unable to deal with the reality of polio, often disappeared (22). It was not uncommon for neighborhood children to be prohibited from playing with "polio victims" for fear that the "crippling" was contagious (23). Polio survivors, no longer physically able to participate easily or fully in social activities, became isolated (10, 25). Such alienation and isolation made polio survivors pariahs (10,14,22,25).

However, there was "hope." Individuals had been imbued with the "unqualified progress ideology" of physical therapy following their acute illness and had often been given prognoses upon discharge that were unrealistically optimistic (26). One paraplegic woman was told, "You'll walk out of your braces before you start dating." Such prognoses were seen as a call to normalcy. The use of braces, crutches and wheelchairs, which had made mobility possible and had been symbols of triumph over paralysis in the hospital, became stigmata of "the crippled" and symbols of polio survivors' failure to have recovered completely. Necessary assistive devices were quickly discarded, regardless of the discomfort, fatigue, awkwardness or pain that resulted, as individuals strove for the "appearance of complete physical normalcy" (21, 27). Painful and exhausting physical therapies were resumed upon patients' return home from hospital or initiated in those who were not hospitalized. Physical therapy often continued for more than a decade after the acute polio, the only goal being "complete recovery" (11). One patient, who became triplegic at age six and has consistently thereafter used a wheelchair, continued daily physical therapy until she left home for college: "Everyone believed that physical therapy would make me walk - eventually." As motivation to walk without braces, another patient was repeatedly beaten by his father who said, "I'd rather see you dead than a cripple".

The abuse of polio survivors by parents was usually less overt but was not infrequent, being motivated by a denial of polio and revulsion by disability (13,14, 27). Some polio survivors report being physically trapped by their parents' refusal to make accommodations for their physical limitations. Homes frequently remained inaccessible, making activities of daily living impossible without assistance from parents - assistance that could be inconsistently provided or consistently withheld (28). One patient was regularly carried outside in the early morning, regardless of the weather, where she remained until dark. Another patient reported that he was locked in a car with windows closed in the heat of summer so that his family could tour Washington's then inaccessible national monuments. Polio survivors learned that physical abilities needed to be maximized if they were to survive in a "barrier-full" society.

Other parents strove to reincorporate the child into the family routine and "forget" about polio by requiring children to equal or exceed the level of physical performance they exhibited before their illness (11). One patient's family took up hiking when their daughter returned home from the hospital, expecting her to "learn to keep up with the others" in spite of crutches and braces. Patients also report that they were expected to out-perform their peers academically. Several patients reported that they would be severely punished if they received "anything less than an 'A'." Such academic expectations reinforced the premature adult-like behaviors that were described in children who had been hospitalized following the acute polio (22).

Still other parents experienced both "shame and estrangement" at having been "singled out" by polio (22). Unable to deny the reality of their child's disability, they isolated the family from society and severely restricted the child's physical activities (11,14,22). One patient, who had a barely noticeable limp following polio at 18 months, was prevented from leaving his yard or playing with other children

until he attended high school. Such isolation impaired the development of social relationships (25,29,30) as parental shame of disability was learned by the child (14).

Unfortunately, many adult patients report that their parents have consistently refused through the years to discuss anything having to do with the physical or emotional reality of the polio experience (14,20). An attempt at discussion with one 45-year old patient's mother was stopped by the angry statement, "It's all too painful for me. Don't ever mention it again." Whether unrealistically demanding or overly restrictive, parents' powerful expectations, combined with those of physicians, nurses and therapists, dictated the behaviors that were required of polio survivors if they were to escape "mortal danger" (15) and be at least partially accepted -- were not totally rejected -- by parents and society. Yet, despite the fulfillment of these expectations by the achievement of a remarkable degree of functional recovery in most individuals and even the appearance of "normalcy" in many, polio survivors were told that they were still not acceptable (31). They report being told without evidence by teachers that no one would ever hire them, by parents that no one would ever marry them, and by physicians that they would never have children (28,32,33). Thus, as polio survivors entered adulthood, they had internalized the expectations of those in authority, which led to self-alienation and actual participation in the "execution of the self" (24). Polio survivors continued to follow the prescription for proper behavior set forth in the "Good Chart" even though they had been told that fulfilling the expectations of others would not result in acceptance by society (34).

**Type A Behavior as a Post-Polio Sequelae.** The majority of polio survivors have succeeded in minimizing the appearance of disability, maximizing independence and "disappearing" into society (28). Most polio survivors have learned to walk after discarding nearly all of their original assistive devices (6). Despite society's negative expectations, polio survivors have had more years of formal education and a larger proportion of them are married as compared to the general disabled as well as nondisabled populations (35). Polio survivors also have a higher level of employment as compared to the general disabled population (36). These data suggested the hypothesis that polio survivors have not merely achieved "normalcy" but have actually surpassed it by learning physically and emotionally stressful Type A behavior as a response to their experience of polio (5,28,37). The polio experience should have provided an ideal environment for the conditioning of Type A behavior. Type A behaviors are thought to be learned in service of the "active avoidance of punishment" (38) by individuals who are engaged in a chronic "struggle to overcome environmental barriers" (39) "against the opposing efforts of other things or persons" (40). Further, Powell, et al (41) found that a "consistent set of beliefs and attitudes about the self, others and life in general lay beneath the overt (Type A) behavior," including "low social support," "low self-esteem" and loss of control. Such beliefs would be the likely consequence of the above-described experience of polio.

To test the hypothesis that polio survivors are Type A, the Brief Type A Questionnaire (42) was administered to 738 polio survivors as part of the 1985 National Post-Polio Survey (5). Polio survivors' mean Type A score of  $53 \pm 22$  was significantly higher than that of non-disabled controls ( $36 \pm 14$ ). This finding was replicated by the 1990 National Survey of 373 polio survivors (6) who had a significantly elevated mean Type A score of  $59 \pm 22$  as compared to nondisabled controls ( $45 \pm 20$ ) and adults with spina bifida ( $48 \pm 22$ ). The 1995 International Survey of 1185 polio survivors again confirmed that polio survivors show significantly more Type A behavior than do non-disabled controls. Averaging all three surveys, polio survivors report 21% more Type A behavior than do those who did not have polio.

Since it had been suggested that Type A individuals are hypersensitive to criticism (41), the 1990 National Survey also included the Reinforcement Motivation Survey (RMS) (43) that contains a Sensitivity to Criticism and Failure scale. Polio survivors had a significantly elevated mean Sensitivity

to Criticism and Failure score ( $68 \pm 28$ ) as compared to non-disabled controls ( $59 \pm 27$ ) that was significantly correlated with the Type A score ( $r=0.44$ )(5). This correlation suggests that Type A behavior in polio survivors serves in part to prevent criticism and to protect against feelings of failure (21,44).

**Compulsive Behavior as a Post-Polio Sequelae.** The majority of patients treated by The Post-Polio Institute not only demonstrate Type A behavior but also evidence behaviors that are not typically Type A. Patients describe an inability to express emotion or admit having physical pain ("I can't complain because people expect the handicapped to complain"). They inflexibly and punitively judge their own behavior on the basis of what is "normal" and on unachievable ideals of perfection. One patient stated, "If I fail at anything, I might as well die. It boils down to either using every ounce of my energy to lead a normal, successful life or giving in to my weaknesses and being inferior."

Patients also demonstrate a strong need to be in control (15) and report marked anxiety with and nearly phobic avoidance of any change ("I am constantly moving and in a constant state of fear. I feel if I slow down I'll never get started again") or decrease in the number or extent of their activities ("I can't just sit and do nothing"). These patients refuse to slow down, delegate responsibilities or allow others to assist them even when they experience fatigue, weakness and pain. They report a saw-tooth pattern of activity, characterized by working until physical symptoms prevent them from continuing. Exhausted, anxious and fearful of criticism, they rest until activity is again possible and then work until symptoms force them to halt. As one polio survivor stated, "I have spent my whole life pushing to keep going and I still push myself even though I know I shouldn't. I keep going until I can no longer walk or stand the pain. I cannot be stopped. I work until I fall."

Patients also report that they "must" satisfy the real or perceived needs of others ("I can't ever say 'No'") and some describe a compulsion to perform nearly ritualistic behaviors in order to escape "mortal danger" (15). One patient stated, "I know it's ridiculous, but I believe that if I make the bed every day I'll never die. I will earn the right to continue to live."

These behaviors are reminiscent of the Obsessive Compulsive Personality Disorder (OCPD)(45). However, polio survivors' compulsive behavior differs from OCPD in important ways. Their compulsivity does not interfere with task completion or promote indecisiveness, nor do polio survivors lack generosity in giving to others. On the contrary, compulsive behaviors are goal-oriented and are performed in the service of others. Importantly, these behaviors are associated with the exacerbation and maintenance of PPS symptoms. Because of these differences with OCPD, a separate diagnostic category -- Compulsive Psychophysiological Disorder (CPD) -- is suggested to highlight the relationship between goal-oriented, compulsive behaviors and the occurrence of physical symptoms.

Fifty percent of the patients evaluated by The Post-Polio Institute met the criteria for CPD. Further, the diagnosis of CPD is significantly correlated with patients' elevated Type A ( $r = .46$ ) and Sensitivity to Criticism and Failure scores ( $r = .47$ ) suggesting that compulsive behaviors are related to Type A behavior and may also serve in part to prevent criticism and to protect against feelings of failure.

These findings support the theoretical literature on compulsive behavior, which suggests that compulsivity protects against loss of control and failure (46) and is used to "avoid or overcome distressful feelings of helplessness" (47) (see "flashbacks," below). Millon and Everly (48) state that compulsive behavior occurs when the "child never develops a separate identity and functions in the world by conforming to strict, internalized parental standards and to the standards around him or her" - conditions that are described above as central to the experience of polio. Additionally, compulsive behavior has been seen to develop in children hospitalized even for brief periods (20).

**Compulsive, Type A Behavior and the Etiology of PPS.** Unfortunately, polio survivors appear to have paid a high price for learning compulsive, Type A behavior in an effort to achieve "normalcy," escape "mortal danger" (15) and protect against the emotional pain generated by the polio experience. That price is Post-Polio Sequelae (28). Physical overexertion and emotional stress were reported to be the first and second leading causes of PPS in the 1985 National Survey (5). The majority of respondents to the 1990 National Survey reported symptoms of chronic stress (frequent feelings of anxiety, low mood and difficulty falling asleep because "my mind is racing")(6,7,8). These symptoms, along with the Type A and Sensitivity to Criticism and Failure scores, were significantly correlated with PPS symptom severity (5,6,49).

These data indicate that PPS are psychophysiological in nature and that clinicians must address both physical and psychological symptoms when polio survivors present for treatment of PPS. Thus, the treatment of PPS requires behavior modification to reduce compulsive, Type A behavior and thereby decrease PPS symptoms. Psychotherapy is also necessary to address the dysfunctional beliefs and powerful suppressed emotions resulting from the experience of polio that motivate compulsive, Type A behavior, produce psychological distress and prevent compliance with therapy.

**Behavioral Modification and Psychotherapy in Treating PPS.** All patients who present for treatment by The Post-Polio Institute undergo complete psychiatric and psychological evaluations, the latter beginning with administration of the Reinforcement Motivation Survey (RMS) (43) and Beck Depression Inventory (BDI) (50). The majority of patients have elevated Type A ( $57 \pm 28$ ) and Sensitivity to Criticism and Failure ( $75 \pm 28$ ) scores on the RMS but had insignificantly elevated BDI scores ( $14 \pm 9$ ).

Although 57% of patients reported "low mood," only 31% met DSM-III-R criteria for Major Depressive Episode (MDE), a percentage similar to the 32% reported by Freidenberg, et al. (51) but higher than the prevalence of MDE in the general population (3% to 7%) (52, 53) and in those with medical illness (20%) (54). Friedenber, et al. (51) suggested that damage to monoaminergic neurons in the brain stem may predispose polio survivors to MDE (see 49). However, the report of "treated depressions" in 38% of adults with spina bifida suggests that childhood-onset orthopedic disability itself may predispose to MDE in adults (55).

The presence of MDE in our patients is important, not only because it requires treatment, but also because MDE is significantly correlated with treatment non-compliance. MDE was diagnosed in 63% of patients who refused further treatment after the initial evaluation and in 50% of those who were discharged for therapeutic non-compliance. Only 11% of patients who were fully compliant and 29% of those who were partially compliant with therapy were diagnosed as having a MDE. MDE is always treated with psychotherapy, although depression was sufficiently severe in 70% of patients that an antidepressant was recommended.

Following psychometric testing, psychological, medical and psychosocial histories are taken that include detailed descriptions of new symptoms and their triggers, the acute polio experience, post-acute polio rehabilitation and patients' personal experience of disability. Information about the response of family, friends and community to the acute polio, to patients' original disability and to PPS is elicited along with a chronology of school and vocational achievements noting the onset and development of compulsive, Type A behavior.

### **Treatment.**

The treatment of PPS is approached by all members of the treatment team - psychiatrist, psychologist and occupational and physical therapists - from a psychophysiological perspective using the techniques of multimodal behavior therapy (56). Therapy begins by asking patients to list their treatment goals and keep a daily log of activities, perceived exertion, fatigue, muscle weakness, pain,

emotional stress, thoughts and emotions. This log is used to document physical and emotional symptoms and demonstrate their relationship to thoughts, affect and compulsive Type A behavior (56). Outpatients are then evaluated by occupational and physical therapists and dieticians during weekly sessions.

After two weeks, log data and evaluation results are used by the treatment team to formulate a behavioral plan designed to decrease behaviors that cause physical symptoms, initiate self-care activities and incorporate stress and time management, energy conservation, work simplification and a program of relaxation and stretching (57) into patients' daily routine. The behavioral plan is then presented to patients and their significant others at a meeting with the treatment team. Also discussed are the hypothesized causes of PPS, the need to modify behavior to treat PPS and the team's expectation of compliance with all aspects of the behavioral plan. Patients continue to keep logs throughout their six to eight weeks of treatment to document symptoms, the progression of physical activity and modification of behaviors. Logs are reviewed with patients by each therapist at every treatment session. Problems with the behavioral plan, especially those having to do with noncompliance, are communicated immediately to all members of the team. Significant problems with compliance may require that patients be called in to meet with the team.

Compliance problems often arise with the daily log and the behavioral plan. Patients have difficulty keeping the log because it interferes with the compulsive performance of their scheduled activities and forces them to recognize the severity and pervasiveness of their symptoms. Invariably, patients have difficulty in complying with the behavioral plan. They will "forget" to alter their schedules and refuse lifestyle modification because such changes directly conflict with their compulsive Type A behavior (46). As one patient who was discharged for non-compliance angrily stated, "I don't want to change. I just want to be the way I was." Unfortunately, such problems with compliance are typical of post-polio patients. One clinic reported that only 41% of patients sporadically used a newly prescribed brace while 70% refused to use a new cane or crutch simply because they "didn't want to" (58). Another clinic concluded that "the major reason for failure (of PPS symptoms) to improve was the unwillingness of the patient to follow any of the recommendations made" (59). Peach (60) has documented the importance of compliance with regard to the treatment of PPS muscle weakness, showing that muscle strength decreases over time only in patients who fail to fully comply with therapeutic recommendations.

On average, 47% of the patients treated by The Post-Polio Institute are fully compliant, 34% partially compliant while 9% are discharged from therapy for not complying with the behavioral plan. Of all patients who received an initial evaluation, 25% refused any treatment for their PPS. It appears that inadequate compliance results in part from patients' fear of relinquishing compulsive, Type A behavior that is, consciously or subconsciously, believed by them to protect against "mortal danger" (15). Some non-compliant patients fear that PPS will return them to the same position of helplessness that they experienced following acute polio (cf. 47). Most report an even greater fear of criticism and a sense of failure when they merely contemplate lifestyle changes or the use of new assistive devices, even though without them PPS symptoms persist, progress and individuals' functional ability deteriorates. Not unexpectedly, inadequately compliant patients have an elevated average Sensitivity to Criticism and Failure score ( $81 \pm 20$ ) as compared to fully compliant patients ( $74 \pm 35$ ).

Patients' ability to change their behavior is directly related to their ability to challenge long-held beliefs about self worth and survival and tolerate the emergence of the powerful fears and long-suppressed emotions generated by the polio experience (41). To promote compliance with the behavioral plan, cognitive therapy techniques are employed through weekly individual psychotherapy to identify and modify dysfunctional beliefs, fears and emotions (46). Learning theory is also employed to explain how compulsive, Type A behavior was learned to protect against fear, emotional pain and "mortal

danger" (15), while behavior modification techniques are used to decrease and eliminate these overlearned behaviors (41,46).

As these techniques assist patients to decrease compulsive Type A behavior, increased anxiety and fear invariably result. Some patients report insomnia, panic attacks, intrusive and emotionally-charged flashbacks about their early polio experience and even fear of impending death (47). The psychotherapist then helps patients understand the origin of these most disturbing experiences and learn to tolerate them, reinforces new self-care behaviors and supports patients through what can be terrifying days and nights.

As therapy continues, patients discover that their compulsive, Type A behavior has been more habitual than necessary. They begin to discard the "Good List," become better able to care for themselves and begin to believe that they can have both self-worth and a disability. As the behaviors and emotions conditioned by the polio experience decrease, the focus of psychotherapy shifts to the experience of PPS. Depressed mood is replaced by sadness and anger as the trauma of a "second disability" is experienced (4). Twice-monthly group psychotherapy provides peer support to assist patients in dealing emotionally with their PPS and practically with the lifestyle changes necessary to treat them. The realization by patients in the group that they are not alone in their painful physical and emotional experiences is itself powerfully therapeutic. Couples and family sessions are also employed for both education about PPS and psychotherapy. It is noteworthy that nearly all spouses, with some striking exceptions, respond favorably to patients' lifestyle changes with the most frequently heard comment being, "I've been wanting you to slow down for years."

All patients treated by The Post-Polio Institute, whether fully or even partially compliant, report meaningful reductions in physical and affective symptoms and increased function (cf. 60). But, as fatigue, muscle weakness and pain decrease, patients frequently increase activity and have a tendency to resume compulsive Type A behavior (46). To prevent this, patients are instructed to strictly adhere to the modified daily schedule as set forth in their behavioral plan, especially after they are discharged from physical and occupational therapies and individual psychotherapy. Patients continue with group psychotherapy for at least six months after discharge, are seen at three-month intervals for medical and psychological follow-up and are encouraged to call their therapists at any time with questions or for support.

## **CONCLUSION**

It is possible that the large proportion of patients treated by The Post-Polio Institute who have experienced severe psychological trauma as a result of their polio experience may not be representative of all polio survivors or even of all individuals who present for treatment of PPS. However, there is something unique about the experience of polio that predisposes to the development of compulsive Type A behavior, sensitivity to criticism and failure, chronic stress, anxiety and depression, i.e., the early onset of a universally feared, potentially life-threatening illness that presented in epidemic proportions; the experience of severe paralysis that could be forced to recede with the application of years of strenuous physical therapy; the need to maximize physical abilities in order to function in a "barrier-full" society that was repelled by disability; the sparing of intellectual abilities that were used to overcompensate for residual physical limitations (61,62). These factors must be studied if we are to understand the psychology of the polio survivor, maximize treatment compliance and effectively treat both the physical and psychological symptoms of PPS.

## REFERENCES

- 1) Gallagher HG: FDR's Splendid Deception. New York, Dodd, Mead, 1985.
- 2) Parsons PE: Data on polio survivors from the National Health Interview Survey. Washington, D.C., National Center for Health Statistics, 1989.
- 3) Bruno RL, Frick NM: Post-Polio Sequelae: Physiological and Psychological Overview. *Rehabilitation Literature* 1986; 47:106-111.
- 4) Frick NM: Post-Polio Sequelae and the psychology of second disability. *Orthopedics* 1985; 8: 851-853.
- 5) Bruno RL, Frick NM: Stress and "Type A" behavior as precipitants of Post-Polio Sequelae, in Halstead LS and Wiechers DO (eds): *Research and CLinical Aspects of the Late Effects of Poliomyelitis*. White Plains, March of Dimes, 1987.
- 6) Bruno RL, Frick NM: Psychological profile of polio survivors. *Arch Phys Med Rehabil* 1990; 10: 889.
- 7) Bruno RL, Frick NM: Five year follow-up to the 1985 National Post-Polio Survey. *Arch Phys Med Rehabil* 1990; 10: 886.
- 8) Conrady LJ, Wish JR, Agre JC, Rodriquez, AA, Sperling, KB: Psychological characteristics of polio survivors. *Arch Phys Med Rehabil* 1989; 70: 458-463.
- 9) Bozarth CH: Unfinished Business: Support Group Members Confront Feelings Surrounding the Late Effects of Polio. Lansing, Polio Survivors Support Group, 1987.
- 10) Copellman FS: Follow-up of one hundred children with poliomyelitis. *The Family* 1944; December: 189-196.
- 11) Meyer E: Psychological considerations in a group of children with poliomyelitis. *J Pediatrics* 1947; 31: 34-48.
- 12) Moustakas CE: *Loneliness*. Englewood Cliffs, Prentice Hall, 1961.
- 13) Seidenfeld MA: The psychological sequelae of poliomyelitis in children. *The Nervous Child* 1948; 7: 14-28.
- 14) Wright BA: *Physical Disability - A psychosocial approach*. New York, Harper and Row, 1983.
- 15) Register C: *Living with Chronic Illness*. New York, Free Press, 1987. 16 Berg RH: *Polio and its Problems*. Philadelphia, Lippincott, 1948.
- 17) Billig HE, van Harrevel A, Wiersma CAG: On re-innervation of paretic muscles by the use of their residual nerve supply. *J Neuropath Exp Neurol* 1946; 5:1-23.
- 18) Kerr N: Staff expectations for disabled persons. *Rehabilitation Counseling Bulletin* 1970; 14: 85-94.
- 19) Hawke WA, Auerbach A: Multidiscipline experience: A fresh approach to aid the multihandicapped child. *Journal of Rehabilitation* 1975; 41:22-24.
- 20) Rowlands P: *Children Apart*. New York, Pantheon, 1973. 21 Schechter MD: The orthopedically handicapped child. *Arch Gen Psychiatry* 1961; 4: 53-59.
- 22) Davis F: *Passage Through Crisis*. Indianapolis, BobbsMerrill, 1963.
- 23) Rice EP: The families of children with poliomyelitis, in *Poliomyelitis*. Philadelphia, Lippincott, 1949.
- 24) Moustakas CE: *Loneliness and Love*. Englewood Cliffs, Prentice Hall, 1972.
- 25) Ladieu-Leviton G, Adler DL, Dembo T: Studies in adjustment to visible injuries: Social acceptance of the injured. *Journal of Social Issues* 1948; 4: 55-62.
- 26) Davis F: *Illness, Interaction and the Self*. Belmont, California, Wadsworth, 1972.
- 27) Long L, Roth C: The psychological aspects of rehabilitation, in Soden WH (ed) *Rehabilitation of the Handicapped*. New York, Ronald, 1949.
- 28) Trieschmann RB: *Aging with a Disability*. New York, Demos, 1987.
- 29) Battle CU: Disruptions in the socialization of a young, severely handicapped child. *Rehabilitation Literature* 1974; 35:130-140.
- 30) Anthony WA: Societal rehabilitation: Changing society's attitudes toward the physically and mentally disabled. *Rehabilitation Psychology* 1972; 19: 117-126.
- 31) Wright BA: Issues in overcoming emotional barriers to adjustment in the handicapped. *Rehabilitation Counseling Bulletin* 1967; 11: 53-59.
- 32) Bloom JL: Sex education for the physically handicapped. *Sexology* 1970; 36: 62-65.
- 33) Hohmann GW: Reactions of the individual with a disability complicated by a sexual problem. *Arch Phys Med Rehabil* 1975; 56: 8-13.
- 34) Ladieu G, Hanfmann E, Dembo T: Studies in adjustment to visible disability. *The Journal of Abnormal Social Psychology* 1947; 42: 169-192.
- 35) Taylor H, Kagay MR, Leichenko S: *The ICD Survey of Disabled Americans*. New York, Louis Harris, 1986.
- 36) *Statistical Abstract of the United States: 1990 (110th edition)*. U.S. Bureau of the Census, Washington, D.C., 1990.
- 37) Bruno RI, Frick NM: The origins of "Type A" behavior and stress induced symptoms in polio survivors. *Proceedings of the Society of Behavioral Medicine* 1989; 10: 85.
- 38) Perkins KA: Heart rate change in Type A and Type B males as a function of response cost and task difficulty. *Psychophysiology* 1984; 21: 14-21.

- 39) Rosenman RH, Friedman M, Straus R, et al: A predictive study of coronary heart disease. *JAMA* 1964; 189: 113-124.
- 40) Friedman M, Thoresen CE, Gill JJ, et al: Feasibility of altering Type A behavior pattern. *Circulation* 1982; 66: 83- 92.
- 41) Powell LH, Friedman M, Thoresen CE, et al: Can the Type A behavior pattern be altered after myocardial infarction? *Psychosom Med* 1984; 46: 293-313.
- 42) Young LD, Barboriak JJ: Reliability of a brief scale for assessment of coronary-prone behavior and standard measures of Type A behavior. *Percept Mot Skills* 1982; 55: 1039-1042.
- 43) Bruno RL: Predicting pain management program outcome: The Reinforcement Motivation Survey. *Arch Phys Med Rehabil* 1990; 10: 785.
- 44) Mandell AJ: Toward a psychobiology of transcendence, in Davidson JM, Davidson RJ (eds): *The Psychobiology of Consciousness*, New York, Plenum, 1980.
- 45) *Diagnostic and Statistical Manual of Mental Disorders* (ed 3, revised). Washington, D.C., American Psychiatric Association, 1987.
- 46) Beck AT, Freeman A: *Cognitive Therapy of Personality Disorders*. New York, Guilford, 1990.
- 47) Salzman L: *The Obsessive Personality*. New York, Jason Aronson, 1973.
- 48) Millon T, Everly G: *Personality and its Disorders*. New York, Wiley, 1985
- 49) Bruno RL, Frick NM, Cohen J. Poliioencephalitis, stress and the etiology of Post-Polio Sequelae. *Orthopedics* 1991; (this issue).
- 50) Beck AT, Ward CH, Mendelson M, et al: An inventory for measuring depression. *Arch Gen Psychiatry* 1961; 4: 561-571.
- 51) Freidenberg DL, Freeman D, Huber SJ, Perry, J, Fischer, A, Van Gorp, WG, et al: Postpoliomyelitis Syndrome: Behavioral Features. *Neuropsychiatry, Neuropsychology and Behavioral Neurology* 1989; 2: 272-281.
- 52) Myers JK, Weissman MM, Tischler OL, et al: Six month prevalence of psychiatric disorders in three communities. *Arch Gen Psychiatry* 1984; 41: 959-967.
- 53) Roberts RE, Vernon SW: Depression in the community: Prevalence and treatment. *Arch Gen Psychiatry* 1982; 39: 1407-1409.
- 54) Schleifer SJ, Macari M, Slater W, et al: Predictors of outcome after myocardial infarction: Role of depression. *Circulation* 1987; 74 (supplement II): 38.
- 55) Dunne KB, Gingham N, Olsen L, et al: The Adult Spina Bifida Network Survey. *Insights* 1986; 14: 28-32.
- 56) Lazarus A: *Multimodal Behavior Therapy*. New York, Springer Verlag, 1976.
- 57) Fillyaw, et al. *Orthopedics* December, 1991.
- 58) Waring WP, Maynard F, Grady W et al: Influence of appropriate lower extremity orthotic management on ambulation, pain and fatigue in a postpolio population. *Arch Phys Med Rehabil* 1989; 70: 371-375.
- 59) Agre JC, Rodriguez AA, Sperling KB: Symptoms and clinical impressions of patients seen in a postpolio clinic. *Arch Phys Med Rehabil* 1989; 70: 367-370.
- 60) Peach PE, Olejnik S: Effects of treatment and noncompliance on post-polio sequelae symptoms and muscle strength. *Orthopedics* 1991; (this issue).
- 61) Scheer J, Luborsky M: Culture and polio. *Orthopedics* 1991; (this issue).